

Exam in Probability Theory, 6 credits

Exam time:	8-12
Allowed:	Pocket calculator. Table with common formulas and moment generating functions (distributed with the exam). Table of integrals (distributed with the exam). Table with distributions from Appendix B in the course book (distributed with the exam).
Examinator:	Mattias Villani.
Assisting teacher:	Per Sidén, phone 0704-977175
Grades:	Grades: Maximum is 20 points. A=19-20 points B=17-18 points C=12-16 points D=10-11 points E=8-9 points F=0-7 points

- Write clear and concise answers to the questions.
 - Make sure to specify the definition region for all density functions.
-

1. The random variable X has the distribution function

$$F_X(x) = \begin{cases} a(1 - \frac{1}{x}), & 1 < x < \infty \\ 0, & \text{otherwise} \end{cases}$$

and the conditional probability density of Y given X is

$$f_{Y|X=x}(y) = \begin{cases} by, & 0 < y < x \\ 0, & \text{otherwise} \end{cases}.$$

- (a) Determine the constant a and the probability density function of X . 1p.
- (b) Determine the constant b as a function of x and compute $E[Y|X=4]$. 1p.
- (c) Compute the joint density function of X and Y . Are X and Y independent? 1p.
- (d) Compute the marginal density of Y and the probability $P(Y < 2)$. 2p.

2. Suppose that X and Y are random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} \frac{2}{3}(2x+y), & \begin{cases} 0 < x < 1 \\ 0 < y < 1 \end{cases} \\ 0, & \text{otherwise} \end{cases}.$$

- (a) Compute $E[2X + Y]$. 2p.
- (b) Determine the distribution of $2X + Y$. 3p.

3. Let X_1 and X_2 follow a multivariate normal distribution with mean vector $\mu = (1, 0)'$ and covariance matrix

$$\Sigma = \begin{pmatrix} 4 & -1 \\ -1 & 1 \end{pmatrix}.$$

Define Y_1, Y_2 and Y_3 through

$$\begin{cases} Y_1 &= X_1 + X_2 \\ Y_2 &= -X_1 + 2X_2 \\ Y_3 &= X_2 - 1 \end{cases}$$

- (a) What is the joint distribution of Y_1, Y_2 and Y_3 ? 1.5p.
(b) Are any of Y_1, Y_2 and Y_3 independent? 1p.
(c) Suppose $X_n \sim \text{Bin}(n, \lambda/n)$. Show that $X_n \xrightarrow{d} \text{Po}(\lambda)$ as $n \rightarrow \infty$. 2.5p.
4. Let $X_k, k = 1, 2, \dots$ be independent random variables, with common density $f_X(x)$ and distribution $F_X(x)$. Also, let N be a positive integer-valued random variable with probability generating function $g_N(t)$. Assume that N and X_1, X_2, \dots are independent. Define

$$Z_N = \max(X_1, X_2, \dots, X_N).$$

- (a) Derive the density of $Z_N|N = n$. 2p.
(b) Show that $F_{Z_N}(z) = g_N(F_X(z))$. 1.5p.
(c) Now, assume X_1, X_2, \dots are all $U(0, 1)$ -distributed and $N \sim \text{Ge}(\frac{1}{2})$. Compute $F_{Z_N}(z)$. 1.5p.

GOOD LUCK!

PER